

CKM Matrix: Status and New Developments

A. Höcker*, H. Lacker*, S. Laplace* and F. Le Diberder*

*Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université de Paris-Sud, BP 34, F-91405 Orsay Cedex, France

Abstract. An analysis of the CKM matrix parameters within the *Rfit* approach is presented using updated input values with special emphasis on the recent $\sin 2\beta$ measurements from BABAR and Belle. The QCD Factorisation Approach describing $B \rightarrow \pi\pi, K\pi$ decays has been implemented in the software package CKMfitter. Fits using branching ratios and CP asymmetries are discussed.

STATISTICAL FRAMEWORK AND INPUTS

In the Standard Model (SM) with three families, CP violation is generated by a single phase in the CKM matrix [1]. This picture can be probed quantitatively by means of a global fit to all quantities sensitive to CKM elements in the SM. The analysis presented here is performed within the *Rfit* statistical approach [2], which is implemented in the software package CKMfitter [3].

The quantity $\chi^2 = -2 \ln \mathcal{L}(y_{\text{mod}})$ is minimized in the fit, where the likelihood function is defined by $\mathcal{L}(y_{\text{mod}}) = \mathcal{L}_{\text{exp}}(x_{\text{exp}} - x_{\text{theo}}(y_{\text{mod}})) \cdot \mathcal{L}_{\text{theo}}(y_{\text{QCD}})$. The experimental part, $\mathcal{L}_{\text{exp}}$, depends on measurements, x_{exp} , and theoretical predictions, x_{theo} , which are functions of model parameters, y_{mod} . The theoretical part, $\mathcal{L}_{\text{theo}}$, describes the knowledge on the QCD parameters, $y_{\text{QCD}} \in \{y_{\text{mod}}\}$, where the theoretical uncertainties are considered as allowed ranges. The agreement between data and the SM is gauged by the global minimum $\chi^2_{\text{min}; y_{\text{mod}}}$, determined by varying all model parameters y_{mod} . For $\chi^2_{\text{min}; y_{\text{mod}}}$, a confidence level (CL), expressing the goodness-of-fit, is computed by means of a Monte Carlo simulation. If the hypothesis “the CKM picture of the SM is correct” is accepted, CLs in parameter subspaces a , e.g. $a = (\bar{\rho}, \bar{\eta})$ [4], are evaluated. For fixed a , one calculates $\Delta\chi^2(a) = \chi^2_{\text{min}; \mu}(a) - \chi^2_{\text{min}; y_{\text{mod}}}$, where μ stands for all model parameters (including y_{QCD}) with the exception of a . The corresponding CL is obtained from $\text{CL}(a) = \text{Prob}(\Delta\chi^2(a), N_{\text{dof}})$, where N_{dof} is the number of degrees of freedom, in general the dimension of the subspace a . Since the CL depends on the choice of the ranges for the y_{QCD} , the results obtained in the fit have to be interpreted with care.

The input values used in this analysis are listed in Tab. 1. For $|V_{ub}|$, inclusive measurements from LEP and exclusive measurements from CLEO have been used. The preliminary CLEO lepton endpoint analysis [5] using moments obtained from $B \rightarrow X_s \gamma$ is not yet included. For $|V_{cb}|$, inclusive measurements from LEP, the measurements of $B \rightarrow D^* \ell \nu$ at zero-recoil and the moments analysis from CLEO [5], using inclusive $B \rightarrow X_c \ell \nu$ and $B \rightarrow X_s \gamma$ decays, have been combined. The uncertainty on Δm_d has been significantly reduced due to the measurements from the *B*-factories [6]. However, the constraint on $(\bar{\rho}, \bar{\eta})$ is not improved since it is dominated by the theoretical uncertainty on $f_{B_d} \sqrt{B_d}$.

TABLE 1. Left: input values for the global fit. Right: fit results quoted for CL > 5 %.

Input Parameter	Value	Fit Output	Range (CL > 5 %)
$ V_{ud} $	0.97394 ± 0.00089	λ	0.2221 ± 0.0041
$ V_{us} $	0.2200 ± 0.0025	A	$0.763 - 0.905$
$ V_{cd} $	0.224 ± 0.014	$\bar{\rho}$	$0.07 - 0.37$
$ V_{cs} $	0.969 ± 0.058	$\bar{\eta}$	$0.26 - 0.49$
$ V_{ub} $	$(3.49 \pm 0.24 \pm 0.55) \cdot 10^{-3}$	$J(10^{-3})$	$2.2 - 3.7$
$ V_{cb} $	$(40.4 \pm 1.3 \pm 0.9) \cdot 10^{-3}$	$\sin 2\alpha$	$-0.90 - 0.51$
Δm_d	$(0.489 \pm 0.008) \text{ ps}^{-1}$	$\sin 2\beta$	$0.59 - 0.88$
Δm_s	Amplitude Spectrum	γ	$37^\circ - 80^\circ$
$ \epsilon_K $	$(2.271 \pm 0.017) \cdot 10^{-3}$	$\Delta m_s (\text{ps}^{-1})$	$14.6 - 32.0$
$\sin 2\beta$	0.793 ± 0.102	$f_{B_d} \sqrt{B_d} (\text{MeV})$	$192 - 284$
$m_t (\overline{\text{MS}})$	$(166 \pm 5) \text{ GeV}$	B_K	$0.52 - 1.68$
$m_c (\text{GeV})$	1.30 ± 0.10	$m_t (\text{GeV})$	$95 - 405$
B_K	$0.87 \pm 0.06 \pm 0.13$	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \cdot (10^{-11})$	$1.6 - 4.4$
η_{cc}	1.38 ± 0.53	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \cdot (10^{-11})$	$4.8 - 9.4$
$f_{B_d} \sqrt{B_d} (\text{GeV})$	$(0.230 \pm 0.028 \pm 0.028) \text{ GeV}$	$\mathcal{B}(B^+ \rightarrow \tau^+ \nu_\tau) \cdot (10^{-7})$	$5.6 - 23.8$
$\eta_B (\overline{\text{MS}})$	0.55 ± 0.01	$\mathcal{B}(B^+ \rightarrow \mu^+ \nu_\mu) \cdot (10^{-5})$	$2.2 - 9.4$
ξ	$1.16 \pm 0.03 \pm 0.05$		




FIGURE 1. Left: constraints in the $(\bar{\rho}, \bar{\eta})$ plane where $\bar{\rho} = \rho(1 - \lambda^2/2)$, $\bar{\eta} = \eta(1 - \lambda^2/2)$. Shown are the 5 % CLs for the individual constraints as shaded areas and the 1 σ - and 2 σ -contours from $\sin 2\beta$. In addition, the 5 % and 90 % CL contours for the combined fit are drawn. Right: constraints in the $(\bar{\rho}, \bar{\eta})$ plane if the likelihood ratio is used for the Δm_s constraint.

For Δm_s , the most recent combined amplitude spectrum from [7] is included in the fit using a modified version of the standard amplitude method [2]. If the amplitude spectrum is translated into a likelihood ratio [8, 9], a stronger constraint is obtained. However, to our knowledge, it has not been demonstrated so far that the likelihood ratio can be inter-




FIGURE 2. Left: Constraints in the $(\bar{\rho}, \bar{\eta})$ plane using R_* (see text) and the bound on R . The CL is indicated by the shaded areas; dark-grey: $> 90\%$, gray: $90\% - 32\%$, light-gray: $32\% - 5\%$. Right: Hypothetical constraints for summer 2002.

interpreted as a probability density function. Hence, we use the more conservative method of Ref. [2] for the numerical analysis presented here. For $\sin 2\beta$, the world average is used. It should be noted that the most precise measurements from BABAR and Belle [6] differ presently by about two standard deviations.

FIT RESULTS

The global minimum of the CKM fit is found to be $\chi^2_{\min, y_{\text{mod}}} = 2.3$, resulting in a goodness-of-fit of 71%. It quantifies the excellent agreement between experimental data and the CKM picture of the SM. Fig. 1 shows the $(\bar{\rho}, \bar{\eta})$ plane. Drawn are 5% CL contours from the single constraints using Δm_d , Δm_s , $|V_{ub}/V_{cb}|$ and $|\epsilon_K|$, respectively, and the 1σ - and 2σ -contours for the four-fold ambiguity on β from $\sin 2\beta$. Shown in addition are the contours for the combined fit including $\sin 2\beta$. The statistical precision of the $\sin 2\beta$ measurement already competes with the indirect, theoretically limited constraints. Fig. 1 (right) illustrates the improved constraint when using the likelihood ratio for Δm_s . Selected numerical results are summarized in Tab. 1.

CHARMLESS TWO BODY DECAYS

Constraints on the angle γ can be obtained from $B \rightarrow \pi\pi, K\pi$ decays. Based on color transparency arguments, theoretical calculations such as the QCD Factorisation Approach (FA) [10] and the QCD hard scattering approach [11] have been developed. Recently, the FA has been implemented in CKMfitter. At present, it is premature to in-

fer reliable constraints on the basis of these calculations due to open theoretical questions [10, 11, 12]. Data from the B -factories are not yet precise enough to probe the calculations in detail. Hence, all fit results are marked by an appropriate “R&D” logo. For this review, a global fit to $B \rightarrow \pi\pi, K\pi$ branching fractions and direct CP asymmetries measured in self-tagging $B \rightarrow K\pi$ decays has been performed within the framework of the FA, where most recent experimental results from BABAR [13], Belle [14] and CLEO [15] have been used. The numerous theoretical parameters are let free to vary within the ranges given in Ref. [10]. We find $\chi^2_{\min; y_{\text{mod}}} = 2.0$ and conclude that data are consistently described within the FA. The best FA fits are found at $\gamma \approx 80^\circ$ and are in agreement with the constraints from the standard fit.

Using less theoretical assumptions, ratios of branching fractions can be formed to derive constraints in the $(\bar{\rho}, \bar{\eta})$ plane. As an example, the CP-averaged ratio

$$R = \frac{\tau_{B^\pm} \mathcal{B}(B^0 \rightarrow K^\pm \pi^\mp)}{\tau_{B^0} \mathcal{B}(B^\pm \rightarrow K^0 \pi^\pm)}, \quad (1)$$

provides the bound $R > \sin^2 \gamma$ which is independent of the strong phases [16]. Unfortunately, the present world average $R = 1.07^{+0.19}_{-0.15}$ leads to weak constraints only owing to the tails of the experimental errors. The ratio

$$R_* = \frac{\mathcal{B}(B^\pm \rightarrow K^0 \pi^\pm)}{2 \cdot \mathcal{B}(B^\pm \rightarrow K^\pm \pi^0)}, \quad (2)$$

measured to be $R_* = 0.70^{+0.16}_{-0.13}$, can be used to derive bounds in the $(\bar{\rho}, \bar{\eta})$ plane [17]. An important input for the theoretical prediction of R_* is the tree-to-penguin ratio (P/T) which can be determined experimentally using the relation

$$\bar{\varepsilon}_{3/2} = R_{\text{th}} \cdot \tan \theta_C \cdot \frac{f_K}{f_\pi} \sqrt{\frac{2 \cdot \mathcal{B}(B^\pm \rightarrow \pi^\pm \pi^0)}{\mathcal{B}(B^\pm \rightarrow K^0 \pi^\pm)}}, \quad (3)$$

where R_{th} stands for SU(3) breaking corrections estimated in the FA to be [10] $R_{\text{th}} = 0.98 \pm 0.05$. The bound on R_* can be translated into a prediction if additional information on the strong phases is inserted [10]. Adopting the values for theoretical ranges quoted in Ref. [10], one obtains the constraints shown in Fig. 2. At present, the constraints remain rather weak due to the limited experimental precision. The slight deformation of the shape pattern around $\gamma \approx 90^\circ$ is due to the bound on R . For summer 2002, an integrated luminosity of 100 fb^{-1} is expected to be collected by each experiment, BABAR and Belle. The experimental precision will then start to provide interesting constraints, as can be seen from Fig. 2 obtained assuming the present central experimental values and appropriately rescaling their errors. However, the constraints would still rely on the validity of some theoretical assumptions not yet fully explored.

Within the FA, the P/T ratio for $B \rightarrow \pi^+ \pi^-$ is predicted. Compared to the present experimental error on the time-dependent asymmetry $S_{\pi\pi} = 0.03^{+0.53}_{-0.56} \pm 0.11$ from BABAR, the quoted theoretical uncertainty is much smaller [10]. In Fig. 3 (left), the constraints in $(\bar{\rho}, \bar{\eta})$ from $S_{\pi\pi}$ are shown using P/T from FA where theoretical uncertainties have been neglected. The right plot shows the constraints when also using $\sin 2\beta$.




FIGURE 3. Left: Constraints on $(\bar{\rho}, \bar{\eta})$ from $S_{\pi\pi}$ using the penguin-over-tree ratio from FA. Right: Constraints when using also $\sin 2\beta$.

ACKNOWLEDGMENTS

It was a pleasure to attend this conference which was very successful despite the tragic events of the 11th September. We are indebted to Martin Beneke and Matthias Neubert for their help implementing the QCD Factorisation Approach in CKMfitter. HL was supported by the Fifth Framework Programme of the European Community Research under the grant No. HPMF-CT-1999-00032.

REFERENCES

1. N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Th. Phys.* **49**, 652 (1973).
2. A. Höcker, H. Lacker, S. Laplace and F. Le Diberder, *Eur. Phys. Jour.* **C21/2**, 225 (2001).
3. “CKMfitter: code, numerical results and plots”, <http://ckmfitter.in2p3.fr/>.
4. L. Wolfenstein, *Phys. Rev. Lett.* **51**, 1945 (1983).
5. R.A. Briere, these proceedings.
6. T. Browder and S. Prell, these proceedings.
7. LEP B Oscillation WG, http://lepobosc.web.cern.ch/LEPBOSC/combined_results/budapest_2001/.
8. H.G. Moser and A. Roussarie, *Nucl. Instr. Meth.* **A384**, 491 (1997).
9. M. Ciuchini *et al.*, *Jour. High Ener. Phys.* **0107**, 013 (2001).
10. M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, hep-ph/0104110, (2001).
11. Y.-Y. Keum, H.-n. Li, and A.I. Sanda, hep-ph/0004173, (2000).
12. M. Ciuchini *et al.*, hep-ph/0104126, (2001).
13. BABAR coll., hep-ex/0105061, (2001); hep-ex/0107074, (2001).
14. Belle coll., hep-ex/0104030 (2001); hep-ex/0106095, (2001).
15. CLEO coll., *Phys. Rev. Lett.* **85**, 515 (2000); *Phys. Rev. Lett.* **85**, 525 (2000).
16. R. Fleischer and T. Mannel, *Phys. Rev.* **D57**, 2752 (1998).
17. M. Neubert and J.R. Rosner, *Phys. Lett.* **B441**, 403 (1998).